

## Bibliographic references

- Töteberg-Harms, M.; Wachtl, J.; Schweier, C.; Funk, J.; Kniestedt, C. Long-term efficacy of combined phacoemulsification plus trabeculectomy versus phacoemulsification plus excimer laser trabeculotomy, *Klinische Monatsblätter für Augenheilkunde (efirst)* DOI: 10.1055/s-0043-100291 *Klinische Studie*, Mar 2017
- Berlin, MD, Michael S.; Töteberg-Harms, MD, Marc; Roka, MD, Vigan; Kleineberg, MD, Lea; Stodtmeister, MD, Richard; Giers, MD, Ulrich; Riggs, BA, Michael E. Excimer Laser Trabeculostomy (ELT), A Laser Based MIGS Procedure With No Implants, Enables Consistent Intraocular Pressure-Lowering In Glaucoma Patients Over 8 Years, Both Stand-Alone And Combined With, *AGS Annual meeting* Mar 2017 ID:113
- J.R. Samples, I.I.K. Ahmed (eds.), *Surgical Innovations in Glaucoma*, DOI 10.1007/978-1-4614-8348-9\_8, © Springer Science+Business Media New York 2014
- Töteberg-Harms M, Hanson JV, Funk J. Cataract surgery combined with excimer laser trabeculotomy to lower intraocular pressure: effectiveness dependent on preoperative IOP. *BMC Ophthalmol.* 2013 Jun 24;13:24. doi: 10.1186/1471-2415-13-24. PubMed PMID: 23799932; PubMed Central PMCID: PMC3724507.
- Berlin MS, Kleinberg L, Stodtmeister R, Spitz J, Giers U. The IOP lowering efficacy of excimer-laser-trabeculostomy in open angle glaucoma patients remains consistent over 5 years. Poster session presented at: 23rd annual meeting of the American Glaucoma Society, San Francisco, 1-3 Mar 2013.
- Stodtmeister R, Kleinberg L, Berlin M, Pillunat L, Giers U. Excimer laser trabeculostomy : 5-year follow-up . Abstract submitted for: the 111th congress of the German Association of Ophthalmologists, Berlin, 19-22 Sept 2013.
- Töteberg-Harms M, et al. One-year results after combined cataract surgery and excimer laser trabeculotomy for elevated intraocular pressure. *Ophthalmologie.* 2011;108(8):733-8.
- Pache M, et al. Laser surgery for glaucoma : excimer-laser trabeculotomy. *Klin Monbl Augenheilkd.* 2006;223(4):303-7.
- Wilmsmeyer S, et al. Excimer laser trabeculotomy: a new, minimally invasive procedure for patients with glaucoma. *Graefes Arch Clin Exp Ophthalmol.* 2006;244:670-6.
- Babighian S, et al. Efficacy and safety of ab interno excimer laser trabeculotomy in primary open-angle glaucoma: two years of follow-up. *Ophthalmologica.* 2006;220:285-90.
- Jampel HD, Musch DC, Gillespie BW, et al., the CIGTS Study Group. Perioperative complications of trabeculectomy in the Collaborative Initial Glaucoma Treatment Study (CIGTS) . *Am J Ophthalmol.* 2005;140:16-22.
- Walker R, Specht H. Theoretical and physical aspects of excimer laser trabeculotomy (ELT) ab interno with the AIDA laser operating at a wavelength of 308 nm. *Biomed Tech (Berl).* 2002; 47:106.
- Neuhann T, Scharrer A, Haefliger E. Excimer laser trabecular ablation ab interno (ELT) in the treatment of chronic openangle glaucoma, a pilot study. *Ophthalmochirurgie.* 2001;13:3.
- Huang S, Yu M, Feng G, Zhang P, Qiu C. Histopathological study of trabeculum after excimer laser trabeculectomy ab internmo . *Yan Ke Xue Bao.* 2001;17(1):11-5.
- Vogel M, Lauritzen K. Selective excimer laser ablation of the trabecular meshwork . *Clinical results. Ophthalmologie.* 1997;94(9): 665-7.
- Lauritzen K, Vogel M. Trabecular meshwork ablation with excimer laser - a new concept of therapy for glaucoma patients. *Invest Ophthalmol Vis Sci.* 1997;38:826.

Jahn R, et al. Macroscopic and microscopic findings after excimer laser treatment of different tissue. J Clin Laser Med Surg. 1992; 10:413.

Kaufmann R, Hibst R. Pulsed Er:YAG and 308 nm UV excimer laser: an in vitro and in vivo study of skin-ablative effects. Laser Surg Med. 1989;9:132.

Berlin MS. Excimer laser applications in glaucoma surgery. Ophthalmol Clin North Am. 1988;1:255.

Berlin MS, Rajacich G, Duffy M. Excimer laser photoablation in glaucoma filtering surgery. Am J Ophthalmol. 1987;103: 713-714.3.

Berlin MS, et al. Excimer laser photoablation in glaucoma filtering surgery. Am J Ophthalmol. 1984;103:713.